

A Method for Computing the Edge-Hyper-Wiener Index of Partial Cubes and an Algorithm for Benzenoid Systems

Niko Tratnik

Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia

Abstract

The edge-hyper-Wiener index of a connected graph G is defined as $WW_e(G) = \frac{1}{2} \sum_{\{e,f\} \subseteq E(G)} d(e,f) + \frac{1}{2} \sum_{\{e,f\} \subseteq E(G)} d(e,f)^2$. We develop a method for computing the edge-hyper-Wiener index of partial cubes, which constitute a large class of graphs with a lot of applications. It is also shown how the method can be applied to trees. Furthermore, an algorithm for computing the edge-hyper-Wiener index of benzenoid systems is obtained. Finally, the algorithm is used to correct already known closed formulas for the edge-Wiener index and the edge-hyper-Wiener index of linear polyacenes.

Keywords: edge-hyper-Wiener index, edge-Wiener index, partial cube, benzenoid system, linear polyacene

2010 MSC: 92E10, 05C12, 05C85, 05C90

1. Introduction

The hyper-Wiener index is a distance-based graph invariant, used as a structure-descriptor for predicting physicochemical properties of organic compounds (often those significant for pharmacology, agriculture, environment-protection etc.). It was introduced in 1993 by M. Randić [1] and has been extensively studied in many papers. Randić's original definition of the hyper-Wiener index was applicable just to trees and therefore, the hyper-Wiener index was later defined for all graphs [2]. The hyper-Wiener index is closely related to the well known Wiener index [3], which is one of the most popular molecular descriptors. Moreover, it is also connected to the Hosoya polynomial [4], since the relationship between them was proved in [5].

In [6] a method for computing the hyper-Wiener index of partial cubes was developed and later it was applied to some chemical graphs [7, 8, 9]. Partial cubes constitute a large class of graphs with a lot of applications and includes, for example, many families of chemical graphs (benzenoid systems, trees, phenylenes, cyclic phenylenes, polyphenylenes).

The Wiener index and the hyper-Wiener index are based on the distances between pairs of vertices in a graph and therefore, similar concepts have been introduced for distances between pairs of edges under the names the edge-Wiener index [10] and the edge-hyper-Wiener index [11], respectively. With other words, the edge-hyper-Wiener index of a graph G is just the hyper-Wiener index of the line graph $L(G)$ and a

Email address: niko.tratnik1@um.si (Niko Tratnik)

similar result holds for the edge-Wiener index. For some recent studies on the edge-hyper-Wiener index see [12, 13, 14, 15].

In this paper we develop a method for computing the edge-hyper-Wiener index of partial cubes. Our main result is parallel to the result from [6]. However, the present method is a bit more difficult and the proof requires some additional insights. We use the cut method to reduce the problem of calculating the edge-hyper-Wiener index of partial cubes to the problem of calculating the edge-Wiener index and the contributions of pairs of Θ -classes (for more information about the cut method see [16]). As an example, we describe how the method can be used on trees. In the case of polycyclic molecules the calculation of the edge-hyper-Wiener index by the definition is not easy, especially if one is interested in finding general expressions of homologous series. Therefore, using the main result and the results from [17], we develop an algorithm for computing the edge-hyper-Wiener index of benzenoid systems, which is much more efficient than the calculation by the definition. Finally, our algorithm is used to obtain closed formulas for the edge-Wiener index and the edge-hyper-Wiener index of linear polyacenes. Although these formulas were previously developed in [18, 11], the results are not correct.

2. Preliminaries

Unless stated otherwise, the graphs considered in this paper are connected. We define $d(x, y)$ to be the usual shortest-path distance between vertices $x, y \in V(G)$.

The *Wiener index* and the *edge-Wiener index* of a connected graph G are defined in the following way:

$$W(G) = \sum_{\{u,v\} \subseteq V(G)} d(u, v), \quad W_e(G) = \sum_{\{e,f\} \subseteq E(G)} d(e, f).$$

The distance $d(e, f)$ between edges e and f of graph G is defined as the distance between vertices e and f in the line graph $L(G)$. Here we follow this convention because in this way the pair $(E(G), d)$ forms a metric space. On the other hand, for edges $e = ab$ and $f = xy$ of a graph G it is also legitimate to set

$$\widehat{d}(e, f) = \min\{d(a, x), d(a, y), d(b, x), d(b, y)\}.$$

Replacing d with $\widehat{d}$, we obtain another variant of the edge-Wiener index:

$$\widehat{W}_e(G) = \sum_{\{e,f\} \subseteq E(G)} \widehat{d}(e, f).$$

Obviously, if e and f are two different edges of G , then $d(e, f) = \widehat{d}(e, f) + 1$. Therefore, for any G it holds

$$W_e(G) = \widehat{W}_e(G) + \binom{|E(G)|}{2}. \quad (1)$$

The *hyper-Wiener index* and the *edge-hyper-Wiener index* of G are defined as:

$$WW(G) = \frac{1}{2} \sum_{\{u,v\} \subseteq V(G)} d(u, v) + \frac{1}{2} \sum_{\{u,v\} \subseteq V(G)} d(u, v)^2,$$

$$WW_e(G) = \frac{1}{2} \sum_{\{e,f\} \subseteq E(G)} d(e,f) + \frac{1}{2} \sum_{\{e,f\} \subseteq E(G)} d(e,f)^2.$$

35 It is easy to see that for any connected graph G it holds $W_e(G) = W(L(G))$ and $WW_e(G) = WW(L(G))$.

The *hypercube* Q_n of dimension n is defined in the following way: all vertices of Q_n are presented as n -tuples $(x_1, x_2, \dots, x_n)$ where $x_i \in \{0, 1\}$ for each $1 \leq i \leq n$ and two vertices of Q_n are adjacent if the corresponding n -tuples differ in precisely one coordinate. Therefore, the *Hamming distance* between two tuples x and y is the number of positions in x and y in which they differ.

40 A subgraph H of a graph G is called an *isometric subgraph* if for each $u, v \in V(H)$ it holds $d_H(u, v) = d_G(u, v)$. Any isometric subgraph of a hypercube is called a *partial cube*. For an edge ab of a graph G , let W_{ab} be the set of vertices of G that are closer to a than to b . We write $\langle S \rangle$ for the subgraph of G induced by $S \subseteq V(G)$. The following theorem puts forth two fundamental characterizations of partial cubes:

Theorem 2.1. [19] *For a connected graph G , the following statements are equivalent:*

- 45 (i) G is a partial cube.
- (ii) G is bipartite, and $\langle W_{ab} \rangle$ and $\langle W_{ba} \rangle$ are convex subgraphs of G for all $ab \in E(G)$.
- (iii) G is bipartite and $\Theta = \Theta^*$.

Is it also known that if G is a partial cube and E is a Θ -class of G , then $G - E$ has exactly two connected components, namely $\langle W_{ab} \rangle$ and $\langle W_{ba} \rangle$, where $ab \in E$. For more information about partial cubes see [19].

50 3. The edge-hyper-Wiener index of partial cubes

In this section a method for computing the edge-hyper-Wiener index of partial cubes is developed. For this purpose, we need some auxiliary results. We start with the following definition.

Definition 3.1. *Let G be a partial cube and let E_k be its Θ -class. Furthermore, let U and U' be the connected components of the graph $G - E_k$. If $e, f \in E(G)$, we define*

$$\delta_k(e, f) = \begin{cases} 1; & e \in E(U) \& f \in E(U') \text{ or } e \in E(U') \& f \in E(U), \\ 0; & \text{otherwise.} \end{cases}$$

55 The following lemma is crucial for our main theorem.

Lemma 3.2. *Let G be a partial cube and let d be the number of its Θ -classes. If $e, f \in E(G)$, then it holds*

$$\widehat{d}(e, f) = \sum_{k=1}^d \delta_k(e, f).$$

Proof. Let a and x be end-vertices of e and f , respectively, such that $d(a, x) = \widehat{d}(e, f)$. Also, let b and y be the remaining end-vertices of e and f , respectively. Furthermore, let P be a shortest path between a and x . Obviously, $|E(P)| = \widehat{d}(e, f)$. If $E_1, \dots, E_d$ are Θ -classes of G , than for every $k \in \{1, \dots, d\}$ we define

$$F_k = E_k \cap E(P).$$

Let $k \in \{1, \dots, d\}$ and let U and U' be the connected components of the graph $G - E_k$ such that $a \in V(U)$. Now we can show that $|F_k| = \delta_k(e, f)$. We consider the following two cases.

1. $F_k = \emptyset$

In this case, since no edge of P is in E_k , it follows that path P is completely contained in U or U' .

60 Hence, $\delta_k(e, f) = 0 = |F_k|$ and the statement is true.

2. $F_k \neq \emptyset$

In this case, since no two edges in a shortest path are in relation Θ , it follows $|F_k| = 1$. Let uv be the edge in F_k . Without loss of generality suppose that $V(U) = W_{uv}$ and $V(U') = W_{vu}$. Obviously, $a \in V(U)$ and $x \in V(U')$. We first show that $b \in V(U) = W_{uv}$. If $b \in W_{vu}$, then $d(b, v) < d(b, u) = d(a, u) + 1$ (the equality holds since G is bipartite) and therefore, $d(b, v) \leq d(a, u)$. Hence, $\widehat{d}(e, f) < |E(P)|$, which
65 is a contradiction. In a similar way we can show that $y \in V(U')$. Therefore, $e \in E(U)$ and $f \in E(U')$. It follows that $\delta_k(e, f) = 1 = |F_k|$.

We have proved that $|F_k| = \delta_k(e, f)$ for any $k \in \{1, \dots, d\}$. Therefore,

$$\widehat{d}(e, f) = |E(P)| = \sum_{k=1}^d |E_k \cap E(P)| = \sum_{k=1}^d |F_k| = \sum_{k=1}^d \delta_k(e, f),$$

which completes the proof. $\square$

To prove the main result, we need to introduce some additional notation. If G is a graph with Θ -classes
70 $E_1, \dots, E_d$, we denote by U_k and U'_k the connected components of the graph $G - E_k$, where $k \in \{1, \dots, d\}$. For any $k, l \in \{1, \dots, d\}$ set

$$\begin{aligned} M_{kl}^{11} &= E(U_k) \cap E(U_l), \\ M_{kl}^{10} &= E(U_k) \cap E(U'_l), \\ M_{kl}^{01} &= E(U'_k) \cap E(U_l), \\ M_{kl}^{00} &= E(U'_k) \cap E(U'_l). \end{aligned}$$

Also, for $k, l \in \{1, \dots, d\}$ and $i, j \in \{0, 1\}$ we define

$$m_{kl}^{ij} = |M_{kl}^{ij}|.$$

Lemma 3.3. *Let G be a partial cube and let d be the number of its Θ -classes. Then*

$$\sum_{e \in E(G)} \sum_{f \in E(G)} \widehat{d}(e, f)^2 = 2\widehat{W}_e(G) + 4 \sum_{k=1}^{d-1} \sum_{l=k+1}^d \left(m_{kl}^{11} m_{kl}^{00} + m_{kl}^{10} m_{kl}^{01} \right).$$

Proof. Using Lemma 3.2 we obtain

$$\begin{aligned}
\sum_{e \in E(G)} \sum_{f \in E(G)} \widehat{d}(e, f)^2 &= \sum_{e \in E(G)} \sum_{f \in E(G)} \left(\sum_{k=1}^d \delta_k(e, f) \right)^2 \\
&= \sum_{e \in E(G)} \sum_{f \in E(G)} \sum_{k=1}^d \sum_{l=1}^d \delta_k(e, f) \delta_l(e, f) \\
&= \sum_{k=1}^d \sum_{l=1}^d \left(\sum_{e \in E(G)} \sum_{f \in E(G)} \delta_k(e, f) \delta_l(e, f) \right) \\
&= \sum_{k=1}^d \sum_{e \in E(G)} \sum_{f \in E(G)} \delta_k(e, f) + \sum_{k=1}^d \sum_{\substack{l=1 \\ l \neq k}}^d \left(\sum_{e \in E(G)} \sum_{f \in E(G)} \delta_k(e, f) \delta_l(e, f) \right) \\
&= \sum_{e \in E(G)} \sum_{f \in E(G)} \sum_{k=1}^d \delta_k(e, f) + 2 \sum_{k=1}^d \sum_{\substack{l=1 \\ l \neq k}}^d \left(\frac{1}{2} \sum_{e \in E(G)} \sum_{f \in E(G)} \delta_k(e, f) \delta_l(e, f) \right) \\
&= \sum_{e \in E(G)} \sum_{f \in E(G)} \widehat{d}(e, f) + 2 \sum_{k=1}^d \sum_{\substack{l=1 \\ l \neq k}}^d \left(m_{kl}^{11} m_{kl}^{00} + m_{kl}^{10} m_{kl}^{01} \right),
\end{aligned}$$

where the last equality follows from the obvious fact that $\delta_k(e, f) \delta_l(e, f) = 1$ if and only if $\delta_k(e, f) = 1$ and $\delta_l(e, f) = 1$. Hence,

$$\sum_{e \in E(G)} \sum_{f \in E(G)} \widehat{d}(e, f)^2 = 2\widehat{W}_e(G) + 4 \sum_{k=1}^{d-1} \sum_{l=k+1}^d \left(m_{kl}^{11} m_{kl}^{00} + m_{kl}^{10} m_{kl}^{01} \right)$$

and we are done. $\square$

Now everything is prepared for the main result of the paper.

Theorem 3.4. *Let G be a partial cube and let d be the number of its Θ -classes. Then*

$$WW_e(G) = 2W_e(G) + \sum_{k=1}^{d-1} \sum_{l=k+1}^d \left(m_{kl}^{11} m_{kl}^{00} + m_{kl}^{10} m_{kl}^{01} \right) - \binom{|E(G)|}{2}. \quad (2)$$

Proof. We first notice that

$$\begin{aligned}
\sum_{e \in E(G)} \sum_{f \in E(G)} d(e, f)^2 &= \sum_{e \in E(G)} \sum_{\substack{f \in E(G) \\ f \neq e}} d(e, f)^2 \\
&= \sum_{e \in E(G)} \sum_{\substack{f \in E(G) \\ f \neq e}} \left(\widehat{d}(e, f) + 1 \right)^2 \\
&= \sum_{e \in E(G)} \sum_{\substack{f \in E(G) \\ f \neq e}} \widehat{d}(e, f)^2 + 2 \sum_{e \in E(G)} \sum_{\substack{f \in E(G) \\ f \neq e}} \widehat{d}(e, f) + \sum_{e \in E(G)} \sum_{\substack{f \in E(G) \\ f \neq e}} 1.
\end{aligned}$$

Using Lemma 3.3 we thus get

$$\sum_{e \in E(G)} \sum_{f \in E(G)} d(e, f)^2 = 2\widehat{W}_e(G) + 4 \sum_{k=1}^{d-1} \sum_{l=k+1}^d \left(m_{kl}^{11} m_{kl}^{00} + m_{kl}^{10} m_{kl}^{01} \right) + 4\widehat{W}_e(G) + |E(G)| \cdot (|E(G)| - 1).$$

Therefore, using also Equation 1, we obtain

$$\begin{aligned}
WW_e(G) &= \frac{1}{4} \sum_{e \in E(G)} \sum_{f \in E(G)} d(e, f) + \frac{1}{4} \sum_{e \in E(G)} \sum_{f \in E(G)} d(e, f)^2 \\
&= \frac{1}{2} W_e(G) + \frac{1}{2} \widehat{W}_e(G) + \sum_{k=1}^{d-1} \sum_{l=k+1}^d \left(m_{kl}^{11} m_{kl}^{00} + m_{kl}^{10} m_{kl}^{01} \right) \\
&\quad + \widehat{W}_e(G) + \frac{|E(G)| \cdot (|E(G)| - 1)}{4} \\
&= 2W_e(G) + \sum_{k=1}^{d-1} \sum_{l=k+1}^d \left(m_{kl}^{11} m_{kl}^{00} + m_{kl}^{10} m_{kl}^{01} \right) - \frac{|E(G)| \cdot (|E(G)| - 1)}{2}
\end{aligned}$$

and the proof is complete. $\square$

As already mentioned, trees are partial cubes. Moreover, a Θ -class in a tree is just a single edge. Therefore, for a tree T with edges $e_1, \dots, e_m$, Theorem 3.4 reduces to

$$WW_e(T) = 2W_e(T) + \sum_{k=1}^{m-1} \sum_{l=k+1}^m m_1(e_k, e_l) m_2(e_k, e_l) - \binom{m}{2},$$

where $m_1(e_k, e_l)$ and $m_2(e_k, e_l)$ are the number of edges in the two extremal connected components of the graph $T - \{e_k, e_l\}$, see Figure 1.




Figure 1: A tree with two extremal components with respect to e_k and e_l .

4. Algorithm for benzenoid systems

Let $\mathcal{H}$ be the hexagonal (graphite) lattice and let Z be a cycle on it. Then a *benzenoid system* is induced by the vertices and edges of $\mathcal{H}$, lying on Z and in its interior. These graphs are the molecular graphs of the benzenoid hydrocarbons, a large class of organic molecules. For more information about benzenoid systems see [20].

An *elementary cut* C of a benzenoid system G is a line segment that starts at the center of a peripheral edge of a benzenoid system G , goes orthogonal to it and ends at the first next peripheral edge of G . By

C we sometimes also denote the set of edges that are intersected by the corresponding elementary cut. Elementary cuts in benzenoid systems have been described and illustrated by numerous examples in several earlier articles.

The main insight for our consideration is that every Θ -class of a benzenoid system G coincide with exactly one of its elementary cuts. Therefore, it is not difficult to check that all benzenoid systems are partial cubes.

In the case of benzenoid systems, Theorem 3.4 provides a particular simple procedure for computing the edge-hyper-Wiener index:

1. We first compute the edge-Wiener index using the procedure described in [17], which we briefly repeat: the edge set of a benzenoid system G can be naturally partitioned into sets E_1, E_2 , and E_3 of edges of the same direction. For $i \in \{1, 2, 3\}$, set $G_i = G - E_i$. Then the connected components of the graph G_i are paths. The quotient graph T_i , $1 \leq i \leq 3$, has these paths as vertices, two such paths (i.e. components of G_i) P' and P'' being adjacent in T_i if some edge in E_i joins a vertex of P' to a vertex of P'' . It is known that T_1, T_2 and T_3 are trees. We next extend the quotient trees T_1, T_2, T_3 to weighted trees $(T_i, w_i), (T_i, w'_i), (T_i, w_i, w'_i)$ as follows:

- for $C \in V(T_i)$, let $w_i(C)$ be the number of edges in the component C of G_i ;
- for $E = C_1C_2 \in E(T_i)$, let $w'_i(E)$ be the number of edges between components C_1 and C_2 .

Then the edge-Wiener index of a benzenoid system G can be computed as

$$W_e(G) = \sum_{i=1}^3 \left(\widehat{W}_e(T_i, w'_i) + W_v(T_i, w_i) + W_{ve}(T_i, w_i, w'_i) \right) + \binom{|E(G)|}{2}. \quad (3)$$

To efficiently compute all the terms in Equation 3, some additional notation is needed. If T is a tree and $e \in E(T)$, then the graph $T - e$ consists of two components that will be denoted by $C_1(e)$ and $C_2(e)$. For a vertex-edge weighted tree (T, w, w') and $e \in E(T)$ set

$$n_i(e) = \sum_{u \in V(C_i(e))} w_i(u) \quad \text{and} \quad m_i(e) = \sum_{e \in E(C_i(e))} w'_i(e).$$

Using this notation we recall the following results (see [17]):

$$\begin{aligned} W_v(T, w) &= \sum_{e \in E(T)} n_1(e)n_2(e), \\ \widehat{W}_e(T, w') &= \sum_{e \in E(T)} m_1(e)m_2(e), \\ W_{ve}(T, w, w') &= \sum_{e \in E(T)} (n_1(e)m_2(e) + n_2(e)m_1(e)). \end{aligned}$$

2. Let us denote the second term in Equation 2 by $WW_e^*(G)$, i.e.

$$WW_e^*(G) = \sum_{k=1}^d \sum_{l=k+1}^d \left(m_{kl}^{11} m_{kl}^{00} + m_{kl}^{10} m_{kl}^{01} \right).$$




Figure 2: Two different positions of two elementary cuts.

To compute this term we proceed as follows. Let C_k and C_l be two distinct elementary cuts (Θ -classes) of a benzenoid system G such that $l > k$. Then there are two different cases, since the elementary cuts can intersect or not - see Figure 2.

By a_{kl} , b_{kl} , c_{kl} , and d_{kl} we denote the number of edges in the corresponding components of $G - C_k - C_l$.

110

Then the contribution of the pair C_k, C_l to $WW_e^*(G)$ will be denoted by $f(C_k, C_l)$ and we obtain

$$f(C_k, C_l) = \begin{cases} a_{kl}b_{kl} + c_{kl}d_{kl}, & C_k \text{ and } C_l \text{ intersect} \\ a_{kl}b_{kl}, & \text{otherwise.} \end{cases}$$

Therefore, term $WW_e^*(G)$ can be computed as

$$WW_e^*(G) = \sum_{k=1}^{d-1} \sum_{l=k+1}^d f(C_k, C_l).$$

Finally, we arrive to the algorithm for the computation of the edge-hyper-Wiener index of benzenoid systems. For a given benzenoid system G we first compute its elementary cuts using a procedure called `calculateElementaryCuts` and then the quotient trees T_i using a procedure `calculateQuotientTrees`. For each T_i we first compute the vertex weights w and the edge weights w' using a procedure `calculateWeights`.

115

When the edge-Wiener index is calculated, we sum up all the contributions of pairs of elementary cuts - procedure `calculateContributionsCuts`. Furthermore, Theorem 3.4 is applied to obtain the final result.

The algorithm thus reads as follows:

Algorithm 1: Edge-Hyper-Wiener Index of Benzenoid Systems

Input : Benzenoid system G with m edges

Output: $WW_e(G)$

```
1  $C_1, \dots, C_d \leftarrow \text{calculateElementaryCuts}(G)$ 
2  $(T_1, T_2, T_3) \leftarrow \text{calculateQuotientTrees}(G)$ 
3 for  $i = 1$  to 3 do
4    $(w_i, w'_i) \leftarrow \text{calculateWeights}(T_i, G)$ 
5    $X_{i,1} \leftarrow W_v(T_i, w_i)$ 
6    $X_{i,2} \leftarrow W_e(T_i, w'_i)$ 
7    $X_{i,3} \leftarrow W_{ve}(T_i, w_i, w'_i)$ 
8    $Y_i \leftarrow X_{i,1} + X_{i,2} + X_{i,3}$ 
9 end
10  $W_e(G) \leftarrow Y_1 + Y_2 + Y_3 + \binom{m}{2}$ 
11  $WW_e^*(G) \leftarrow 0$ 
12 for  $k = 1$  to  $d - 1$  do
13   for  $l = k + 1$  to  $d$  do
14      $f_{k,l} \leftarrow \text{calculateContributionsCuts}(C_k, C_l)$ 
15      $WW_e^*(G) \leftarrow WW_e^*(G) + f_{k,l}$ 
16   end
17 end
18  $WW_e(G) \leftarrow 2W_e(G) + WW_e^*(G) - \binom{m}{2}$ 
```

5. Closed formulas for linear polyacenes

120 In this section we derive closed formulas for the edge-Wiener index and the edge-hyper-Wiener index of linear polyacenes using the procedure described in Section 4. If $h \geq 1$, then *linear polyacene* L_h is formed of h linearly connected hexagons, see Figure 3. Recall that for $h = 1, 2, 3, 4, 5$ the graph L_h represents benzene, naphthalene, anthracene, naphthacene, and pentacene (see [20]).




Figure 3: Linear polyacene L_4 .

Although closed formulas for linear polyacenes were previously obtained in [18, 11], the results are not correct.

125 For example, it can be computed by hand that $W_e(L_3) = 350$ and $WW_e(L_3) = 812$, but these results do not

coincide with the mentioned results.

To obtain closed formulas, we first compute the edge-Wiener index. The corresponding weighted quotient trees for L_h are depicted in Figure 4.




Figure 4: Weighted quotient trees for linear polyacene L_h .

Obviously,

$$W_v(T_1, w_1, w'_1) = 4h^2, \quad \widehat{W}_e(T_1, w_1, w'_1) = 0, \quad W_{ve}(T_1, w_1, w'_1) = 0.$$

130 To compute the corresponding Wiener indices of the second tree, we first notice that for the edge e_i , $i \in \{1, \dots, h\}$ it holds:

$$\begin{aligned} n_1(e_i) &= 3(i-1) + 2 = 3i - 1, \\ n_2(e_i) &= 3(h-i) + 2 = 3h - 3i + 2, \\ m_1(e_i) &= 2(i-1) = 2i - 2, \\ m_2(e_i) &= 2(h-i) = 2h - 2i. \end{aligned}$$

Therefore, after an elementary calculation we obtain

$$\begin{aligned} W_v(T_2, w_2, w'_2) = W_v(T_3, w_3, w'_3) &= \sum_{i=1}^h (3i-1)(3h-3i+2) = \frac{1}{2}h(3h^2 + 3h + 2), \\ \widehat{W}_e(T_2, w_2, w'_2) = \widehat{W}_e(T_3, w_3, w'_3) &= \sum_{i=1}^h (2i-2)(2h-2i) = \frac{2}{3}h(h^2 - 3h + 2), \\ W_{ve}(T_2, w_2, w'_2) = W_{ve}(T_3, w_3, w'_3) &= \sum_{i=1}^h \left((3i-1)(2h-2i) + (2i-2)(3h-3i+2) \right) \\ &= 2(h-1)h^2. \end{aligned}$$

Also, one can easily see the the number of edges in L_h is

$$|E(L_h)| = 5h + 1. \quad (4)$$

Hence, Equation 3 imply that for any $h \geq 1$ it holds

$$W_e(L_h) = \frac{1}{6}h(50h^2 + 69h + 43). \quad (5)$$

Next, we have to compute $WW_e^*(L_h)$. Let $B, C_1, C_2, \dots, C_h, D_1, D_2, \dots, D_h$ be the elementary cuts of L_h , where $h \geq 1$. See Figure 5.




Figure 5: Elementary cuts of L_h .

To obtain the final result, we need to calculate the contributions $f(\cdot, \cdot)$ of all the pairs of elementary cuts. Therefore, the number of edges in specific components (parts) of a graph is shown in Table 1.

Pair of elementary cuts	Part a	Part b	Part c	Part d
B, C_k ($k = 1, \dots, h$)	$2k - 1$	$2h - 2k + 1$	$2h - 2k$	$2k - 2$
B, D_k ($k = 1, \dots, h$)	$2k - 2$	$2h - 2k$	$2h - 2k + 1$	$2k - 1$
C_k, C_l ($k < l$)	$5k - 3$	$5h - 5l + 2$	/	/
D_k, D_l ($k < l$)	$5k - 3$	$5h - 5l + 2$	/	/
C_k, D_l ($k < l$)	$5k - 3$	$5h - 5l + 2$	/	/
C_k, D_l ($k > l$)	$5l - 3$	$5h - 5k + 2$	/	/
C_k, D_k ($k = 1, \dots, h$)	$5k - 4$	$5h - 5k + 1$	0	0

Table 1: Number of edges in the connected components.

After some elementary calculations we obtain

$$\begin{aligned}
 \sum_{k=1}^h \left((2k-1)(2h-2k+1) + (2h-2k)(2k-2) \right) &= \frac{1}{3}h(4h^2 - 6h + 5), \\
 \sum_{k=1}^{h-1} \sum_{l=k+1}^h (5k-3)(5h-5l+2) &= \frac{1}{24}h(25h^3 - 70h^2 + 83h - 38), \\
 \sum_{k=1}^h (5k-4)(5h-5k+1) &= \frac{1}{6}h(25h^2 - 45h + 26).
 \end{aligned}$$

By summing up the contributions of all pairs of elementary cuts from Table 1 we then calculate

$$WW_e^*(L_h) = \frac{1}{6}h(25h^3 - 29h^2 + 14h + 8). \quad (6)$$

Finally, using Theorem 3.4, Equation 5, Equation 6, and Equation 4 we deduce that for any $h \geq 1$ it holds

$$WW_e(L_h) = \frac{1}{6}h(25h^3 + 71h^2 + 77h + 79).$$

References

- [1] M. Randić, Novel molecular descriptor for structure-property studies, *Chem. Phys. Lett.* 211 (1993) 478–483.
- [2] D. J. Klein, I. Lukovits, I. Gutman, On the definition of hyper-Wiener index for cycle-containing structures, *J. Chem. Inf. Comput. Phys. Chem. Sci.* 35 (1995) 50–52.
- [3] H. Wiener, Structural determination of paraffin boiling points, *J. Amer. Chem. Soc.* 69 (1947) 17–20.
- [4] H. Hosoya, On some counting polynomials in chemistry, *Discrete Appl. Math.* 19 (1988) 239–257.
- [5] G. G. Cash, Relationship between the hosoya polynomial and the hyper-wiener index, *Appl. Math. Lett.* 15 (2002) 893–895.
- [6] S. Klavžar, Applications of isometric embeddings to chemical graphs, *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.* 51 (2000) 249–259.
- [7] G. Cash, S. Klavžar, M. Petkovšek, Three methods for calculation of the hyper-Wiener index of molecular graphs, *J. Chem. Inf. Comput. Sci.* 42 (2002) 571–576.
- [8] S. Klavžar, P. Žigert, I. Gutman, An algorithm for the calculation of the hyper-Wiener index of benzenoid hydrocarbons, *Comput. Chem.* 24 (2000) 229–233.
- [9] P. Žigert, S. Klavžar, I. Gutman, Calculating the hyper-Wiener index of benzenoid hydrocarbons, *ACH - Models Chem.* 137 (2000) 83–94.
- [10] A. Iranmanesh, I. Gutman, O. Khormali, A. Mahmiani, The edge versions of Wiener index, *MATCH Commun. Math. Comput. Chem.* 61 (2009) 663–672.
- [11] A. Iranmanesh, A. S. Kafrani, O. Khormali, A new version of hyper-Wiener index, *MATCH Commun. Math. Comput. Chem.* 65 (2011) 113–122.
- [12] N. Tratnik, P. Žigert Pleteršek, Relationship between the Hosoya polynomial and the edge-Hosoya polynomial of trees, preprint.
- [13] M. Azari, A. Iranmanesh, Edge-Wiener type invariants of splices and links of graphs, *U.P.B. Sci. Bull., Series A* 77 (2015) 143–154.
- [14] A. Soltani, A. Iranmanesh, The hyper edge-Wiener index of corona product of graphs, *Trans. Comb.* 4 (2015) 1–9.
- [15] A. Soltani, A. Iranmanesh, Z. A. Majid, The multiplicative version of the edge Wiener index, *MATCH Commun. Math. Comput. Chem.* 71 (2014) 407–416.

- 170 [16] S. Klavžar, M. J. Nadjafi-Arani, Cut method: update on recent developments and equivalence of independent approaches, *Curr. Org. Chem.* 19 (2015) 348–358.
- [17] A. Kelenc, S. Klavžar, N. Tratnik, The edge-Wiener index of benzenoid systems in linear time, *MATCH Commun. Math. Comput. Chem.* 74 (2015) 521–532.
- [18] O. Khormali, A. Iranmanesh, I. Gutman, A. Ahmadi, Generalized Schultz index and its edge versions, 175 *MATCH Commun. Math. Comput. Chem.* 64 (2010) 783–798.
- [19] R. Hammack, W. Imrich, S. Klavžar, *Handbook of Product Graphs*, Second Edition, RC Press, Taylor & Francis Group, Boca Raton, 2011.
- [20] I. Gutman, S. J. Cyvin, *Introduction to the Theory of Benzenoid Hydrocarbons*, Springer-Verlag, Berlin, 1989.